

DEPTH AND STANLEY DEPTH OF THE CANONICAL FORM OF A FACTOR OF MONOMIAL IDEALS

ADRIAN POPESCU

ABSTRACT. In this paper we show that the depth and the Stanley depth of the factor of two monomial ideals is invariant under taking a so called canonical form. It follows easily that the Stanley Conjecture holds for the factor if and only if it holds for its canonical form. In particular, we construct an algorithm which simplifies the depth computation and using the canonical form we massively reduce the run time for the sdepth computation.

INTRODUCTION

Let K be a field and $S = K[x_1, \dots, x_n]$ be the polynomial ring over K in n variables. A Stanley decomposition of a graded S -module M is a finite family

$$\mathcal{D} = (S_i, u_i)_{i \in I}$$

in which u_i are homogeneous elements of M and S_i are graded K -algebra retract of S for all $i \in I$ such that $S_i \cap \text{Ann}(u_i) = 0$ and

$$M = \bigoplus_{i \in I} S_i u_i$$

as a graded K -vector space. The Stanley depth of $\mathcal{D}$, denoted by $\text{sdepth}(\mathcal{D})$, is the depth of the S -module $\bigoplus_{i \in I} S_i u_i$. The Stanley depth of M is defined as

$$\text{sdepth}(M) := \max\{\text{sdepth}(\mathcal{D}) \mid \mathcal{D} \text{ is a Stanley decomposition of } M\}.$$

Another definition of sdepth using partitions is given in [4].

Stanley's Conjecture [12] states that the Stanley depth $\text{sdepth}(M)$ is $\geq \text{depth}(M)$.

Let $J \subsetneq I \subset S$ be two monomial ideals in S . In [5], Ichim et. al. studied the sdepth and depth of the factor I/J under polarization and reduced the Stanley's Conjecture to the case when the ideals are monomial squarefree. This is possible the best result from the last years concerning Stanley's depth. It is worth to mention that this result is not very useful for computing sdepth since it introduces a lot of new variables. In the squarefree case there are not many known results about the Stanley conjecture (see for example [9]).

The support from the Department of Mathematics of the University of Kaiserslautern is gratefully acknowledged.

Another result of [5] which seems to help in the computing of sdepth is the following proposition, which extends [2, Lemma 1.1], [6, Lemma 2.1].

Proposition 0.1. [5, Proposition 5.1] *Let $k \in \mathbb{N}$ and I'', J'' be the monomial ideals obtained from I, J in the following way: Each generator whose degree in x_n is at least k is multiplied by x_n and all other generators are taken unchanged. Then $\text{sdepth}_S I/J = \text{sdepth}_S I''/J''$.*

Inspired by this proposition we introduced a canonical form of a factor I/J of monomial ideals (see Definition 1.3) and we showed easily that sdepth is invariant under taking the canonical form (see our Theorem 1.7). This leads us to the idea to study also the depth case (see Theorem 1.11). Our Theorem 1.12 says that Stanley's Conjecture holds for a factor of monomial ideals if and only if it holds for its canonical form. As a side result, in the depth (respectively sdepth) computation algorithm for I/J , one can first compute the canonical form and use the algorithm on this new much more simpler module (see the Appendix).

In Example 1.13 we conclude that the depth and sdepth algorithms are faster when considering the canonical form: using CoCoA[1], SINGULAR[3] and Rinaldo's sdepth computation algorithm [11] we see a small decrease in the depth case timing, but in the sdepth case the run time is massively reduced. We hope that our algorithm together with the one from [8] will be used very often in problems concerning monomial ideals.

We owe thanks to Y.-H. Shen who noticed our results in a previous arXiv version and showed us the papers of Okazaki and Yanagawa [7] and [13], because they are strongly connected with our topic. Indeed Proposition 0.1 and Corollary 1.10 follow from [7, Theorem 5.2] (see also [7, Section 2,3]). However, our proofs of Lemma 1.9 and Corollary 1.10 are completely different from those appeared in the quoted papers and we keep them for the sake of our completeness.

1. THE CANONICAL FORM OF A FACTOR OF MONOMIAL IDEALS

Let $R = K[x_1, \dots, x_{n-1}]$ be the polynomial K -algebra over a field K and $S := R[x_n]$. Consider $J \subsetneq I \subset R$ two monomial ideals and denote by $G(I)$, respectively $G(J)$, the minimal (monomial) system of generators of I , respectively J .

Definition 1.1. The power x_n^r enters in a monomial u if $x_n^r | u$ but $x_n^{r+1} \nmid u$.

We say that I is of type $(k_1, \dots, k_s)$ with respect to x_n if $x_n^{k_i}$ are all the powers of x_n which enter in a monomial of $G(I)$ for $i \in [s]$ and $1 \leq k_1 < \dots < k_s$.

I is in the canonical form with respect to x_n if I is of type $(1, \dots, s)$ for some $s \in \mathbb{N}$.

We simply say that I is the canonical form if it is in the canonical form with respect to all variables $x_1, \dots, x_n$.

Remark 1.2. Suppose that I is of type $(k_1, \dots, k_s)$ with respect to x_n . It is easy to get the canonical form I' of I with respect to x_n : replace $x_n^{k_i}$ by x_n^i whenever

$x_n^{k_i}$ enters in a generators of $G(I)$. Applying by recurrence this procedure for other variables we get the *canonical form* of I , that is with respect to all variables. Note that a squarefree monomial ideal is of type (1) with respect to each x_i and it is in the canonical form with respect to x_i , so in this case $I' = I$.

Definition 1.3. Consider now $J \subsetneq I \subset S$ two monomial ideals. We say that I/J is of type $(k_1, \dots, k_s)$ with respect to x_n if $x_n^{k_i}$ are all the powers of x_n which enter in a monomial of $G(I) \cup G(J)$ for $i \in [s]$ and $1 \leq k_1 < \dots < k_s$. All the terminology presented in Definition 1.1 will extend automatically to the factor case. Thus we may speak about the *canonical form* $\overline{I/J}$ of I/J .

Remark 1.4. In order to compute the canonical form with respect to x_n of the $(k_1, \dots, k_s)$ -type factor I/J , one will replace $x_n^{k_i}$ by x_n^i whenever $x_n^{k_i}$ enters a generator of $G(I) \cup G(J)$.

Example 1.5. We present some examples where we compute the canonical form of a monomial ideal, respectively a factor of two monomial ideals.

- (1) Consider $S = \mathbb{Q}[x, y]$ and the monomial ideal $I = (x^4, x^3y^7)$. Then the canonical form of I is $I' = (x^2, xy)$.
- (2) Consider $S = \mathbb{Q}[x, y, z]$, $I = (x^{10}y^5, x^4yz^7, z^7y^3)$ and $J = (x^{10}y^{20}z^2, x^3y^4z^{13}, x^9y^2z^7)$.

$$\text{The canonical form of } I/J \text{ is } \overline{I/J} = \frac{(x^4y^5, x^2yz^2, y^3z^2)}{(x^4y^6z, xy^4z^3, x^3y^2z^2)}.$$

The canonical form of a factor of monomial ideals I/J is not usually the factor of the canonical forms of I and J as shows the following example.

Example 1.6. Let $S = \mathbb{Q}[x, y]$, $I = (x^4, y^{10}, x^2y^7)$ be and $J = (x^{20}, y^{30})$. The canonical form of I is $I' = (x^2, y^2, xy)$ and the canonical form of J is $J' = (x, y)$. Then $J' \not\subset I'$. But the canonical form of the factor I/J is $\overline{I/J} = \frac{(x^2, y^2, xy)}{(x^3, y^3)}$.

Using Proposition 0.1, we see that the Stanley depth of a monomial ideal does not change when considering its canonical form.

Theorem 1.7. Let I, J be monomial ideals in S and $\overline{I/J}$ the canonical form of I/J . Then

$$\text{sdepth}_S I/J = \text{sdepth}_S \overline{I/J}.$$

The proof goes applying inductively the following lemma.

Lemma 1.8. Suppose that I/J is of type $(k_1, \dots, k_s)$ with respect to x_n and $k_j + 1 < k_{j+1}$ for some $0 \leq j < s$ (we set $k_0 = 0$). Let $G(I')$ (resp. $G(J')$) be the set of monomials obtained from $G(I)$ (resp. $G(J)$) by substituting $x_n^{k_i}$ by $x_n^{k_i-1}$ for $i > j$ whenever $x_n^{k_i}$ enters in a monomial of $G(I)$ (resp. $G(J)$). Let I' and J' be the ideals generated by $G(I')$ and $G(J')$. Then

$$\text{sdepth}_S I/J = \text{sdepth}_S I'/J'.$$

The proof of Lemma 1.8 follows from the proof of [5, Proposition 5.1] (see here Proposition 0.1).

Next we focus on the depth I/J and depth $\overline{I/J}$. The idea of the proof of the following lemma is taken from [10, Section 2].

Lemma 1.9. *Let $I_0 \subset I_1 \subset \dots \subset I_e \subset R$, $J \subset S$, $U_0 \subset U_1 \subset \dots \subset U_e \subset R$, $V \subset S$ be some graded ideals of S , respectively R , such that $U_i \subset I_i$ for $0 \leq i \leq e$, $I_e \subset J$, $V \subset J$ and $U_e \subset V$. Consider $T_k = \sum_{i=0}^e x_n^i I_i S + x_n^k J$ and $W_k = \sum_{i=0}^e x_n^i U_i S + x_n^k V$ for $k > e$. Then $\text{depth}_S \frac{T_k}{W_k}$ is constant for all $k > e$.*

Proof. Consider the following linear subspaces of S : $I := \sum_{i=0}^e x_n^i I_i$ and $U := \sum_{i=0}^e x_n^i U_i$.

Note that I and U are not ideals in S .

If $I = U$, then the claim follows easily from the next chain of isomorphisms $\frac{T_k}{W_k} \cong \frac{x_n^k J}{x_n^k J \cap (I + x_n^k V)S} \cong \frac{x_n^k J}{x_n^k (I + V)S} \cong \frac{J}{(I + V)S}$ for all $k > e$, and hence $\text{depth}_S \frac{T_k}{W_k}$ is constant for all $k > e$.

Assume now that $I \neq U$ and consider the following exact sequence

$$0 \rightarrow \frac{J}{V} \xrightarrow{\cdot x_n^k} \frac{T_k}{W_k} \rightarrow \frac{T_k}{W_k + x_n^k J} \rightarrow 0,$$

where the last term we denote by H_k . Note that $H_k \cong \frac{IS}{IS \cap (U + x_n^k J)S}$ and $IS \cap (U + x_n^k J)S = US + x_n^k IS$. Since $x_n^k H_k = 0$, H_k is a $S/(x_n^k)$ -module. Then $\text{depth}_S H_k = \text{depth}_{S/(x_n^k)} H_k = \text{depth}_R H_k$ because the graded maximal ideal m of R generates a zero dimensional ideal in $S/(x_n^k)$. But H_k over R is isomorphic with $\frac{\bigoplus_{i=0}^{k-1} I_i}{\bigoplus_{i=0}^{k-1} U_i} \cong \bigoplus_{i=0}^{k-1} \frac{I_i}{U_i}$, where $I_i = I_e$ and $U_i = U_e$ for $e < i < k$. It follows that

$$t := \text{depth}_S H_k = \min_i \left\{ \text{depth}_R \frac{I_i}{U_i} \right\}.$$

If $\text{depth}_S \frac{J}{V} = 0$, then the Depth Lemma gives us $\text{depth}_S \frac{T_k}{W_k} = t = 0$ for all $k > e$ and hence we are done. Therefore we may suppose that $\text{depth}_S \frac{J}{V} > 0$.

Note that $t > 0$ implies $\text{depth}_S \frac{T_k}{W_k} > 0$ by the Depth Lemma since otherwise $\text{depth}_S \frac{T_k}{W_k} = \text{depth}_S \frac{J}{V} = 0$, which is false. Next we will split the proof in two cases.

◦ Case $t = 0$.

Let $\mathcal{F} = \{i \in \{0, \dots, e\} \mid \text{depth}_R I_i/U_i = 0\}$ and $L_i \subset I_i$ be the graded ideal containing U_i such that $L_i/U_i \cong H_m^0(I_i/U_i)$.

If $i \in \mathcal{F}$ and there exists $u \in (L \cap V) \setminus U_i$ then $(m^s, x_n^k)x_n^i u \subset W_k$ for some $s \in \mathbb{N}$, that is $\text{depth}_S \frac{T_k}{W_k} = 0$ for all $k > e$.

Now consider the case when $L_i \cap V = U_i$ for all $i \in \mathcal{F}$. If $i \in \mathcal{F}$ then note that $L_i \subset L_j$ for $i < j \leq e$. Set $V' = V + L_e S$, $U' = U + \sum_{i \in \mathcal{F}} x_n^i L_i$ and $W'_k := U' S + x_n^k V' = U' S + x_n^k V$ because $x_n^k L_e S \subset U' S$. Consider the following exact sequence

$$0 \rightarrow \frac{W'_k}{W_k} \rightarrow \frac{T_k}{W_k} \rightarrow \frac{T_k}{W'_k} \rightarrow 0.$$

For the last term we have $H_m^0(I_j/U'_j) = 0$, $0 \leq j \leq e$ and so the new $t > 0$, which is our next case. Thus we get $\text{depth}_S \frac{T_k}{W'_k} > 0$ is constant for $k > e$. The first term

is isomorphic to $\frac{U' S}{U' S \cap W_k}$. But $U' S \cap W_k = U S + (U' S \cap x_n^k V)$ since $U S \subset U' S$. Since $U' S \cap (x_n^k S) = x_n^k (U_e + L_e) S$ and $U_e \subset V$ it follows that $U' S \cap x_n^k V = x_n^k U S + (x_n^k L_e S \cap x_n^k V S) = x_n^k U S$. Consequently, the first term from the above exact sequence is isomorphic with $\frac{U' S}{U S}$. Note that the annihilator of the element

induced by some $u \in L_e \setminus V$ in $U' S / U S$ contains a power of m and so $\text{depth}_S \frac{U' S}{U S} \leq 1$. The inequality is equality since x_n is regular on $U' S / U S$. By the Depth Lemma we get $\text{depth}_S \frac{T_k}{W_k} = 1$ for all $k > e$.

◦ Case $t > 0$.

If $\text{depth}_R \frac{J}{V} \leq t = \text{depth}_S H_k$ then the Depth Lemma gives us again the claim, i.e. $\text{depth}_S \frac{T_k}{W_k} = \text{depth}_S \frac{J}{V}$ for all $k > e$.

Assume that $\text{depth}_S \frac{J}{V} > t$. Apply induction on t , the initial step $t = 0$ being done in the first case. Suppose that $t > 0$. Then $\text{depth}_S \frac{J}{V} > t > 0$ implies that $\text{depth}_S \frac{J}{V} \geq 2$ and so we may find a homogeneous polynomial $f \in m$ that is regular on $\frac{J}{V}$. Moreover we may find f to be regular also on all $\frac{I_i}{U_i}$, $i \leq e$.

Then f is regular on $\frac{T_k}{W_k}$. Set $V'' := V + fJ$ and $U''_i := U_i + fI_i$ for all $i \leq e$

and set $W''_k := \sum_{i=0}^e x_n^i U''_i S + x_n^k V''$. By Nakayama's Lemma we get $U'' \neq U$, and

therefore $\text{depth}_R \frac{I}{U''} = t - 1$ and by induction hypothesis it results that $\text{depth}_S \frac{T_k}{W_k} = 1 + \text{depth}_S \frac{T_k}{W''_k} = \text{constant}$ for all $k > e$.

Finally, note that we may pass from the first case to the second one and conversely. In this way U increases at each step. By Noetherianity at last we may arrive in finite steps to the case $I = U$, which was solved at the beginning. $\square$

The next corollary is in fact [5, Proposition 5.1] (see Proposition 0.1) for depth. It follows easily from Lemma 1.9 but also from [7, Proposition 5.2] (see also [13, Sections 2, 3]).

Corollary 1.10. *Let $e \in \mathbb{N}$, I and J monomial ideals in $S := K[x_1, \dots, x_n]$. Consider I' and J' be the monomial ideals obtained from I and J in the following way: each generator whose degree in $x_n \geq e$ is multiplied by x_n and all the other generators are left unchanged. Then*

$$\text{depth}_S I/J = \text{depth}_S I'/J'.$$

This leads us to the equivalent result of Theorem 1.7 for depth.

Theorem 1.11. *Let I and J be two monomial ideals in S and $\overline{I/J}$ the canonical form of I/J . Then*

$$\text{depth}_S I/J = \text{depth}_S \overline{I/J}.$$

Proof. Assume that I/J is of type $(k_1, \dots, k_s)$ with respect to x_n and obviously $\overline{I/J}$ is of type $(1, 2, \dots, s)$ with respect to x_n . Starting with $\overline{I/J}$, we apply Corollary 1.10 till we obtain an I'_1/J'_1 of type $(k_1, k_1 + 1, \dots, k_1 + s - 1)$ having the same depth as $\overline{I/J}$. We repeat the process until we get I'_s/J'_s of type $(k_1, k_2, \dots, k_s)$ with respect to x_n with the unchanged depth. Now we iterate and take the next variable. At the very end the claim will follow. $\square$

Theorem 1.7 and Theorem 1.11 give us the following theorem

Theorem 1.12. *The Stanley conjecture holds for a factor of monomial ideals I/J if and only if it holds for its canonical form $\overline{I/J}$.*

Using Theorem 1.11, instead of computing the depth or the sdepth of I/J , $J \subsetneq I \subset S$, we can compute it for the simpler module $\overline{I/J}$.

Example 1.13. We present the different timings for the depth and sdepth computation algorithms with and without extracting the canonical form. SINGULAR[3] was used in the depth computations while CoCoA [1] and Rinaldo's paper[11] were used for the Stanley depth computation.

- (1) Consider the ideals from Example 1.5(2).
 Timing for sdepth I/J computation: 22s.
 Timing for sdepth $\overline{I/J}$ computation: 74 ms.
- (2) Consider $R = \mathbb{Q}[x, y, z]$ and $I = (x^{100}yz, x^{50}yz^{50}, x^{50}y^{50}z)$. Then the canonical form is $I' = (x^2yz, xyz^2, xy^2z)$.
 Timing for sdepth I computation: 13m 3s.
 Timing for sdepth I' computation: 21 ms.
 Notice that the difference in timings is very large. Therefore using the canonical form in the sdepth computation is a very important optimization

step. On the other side, the depth computation is immediate in both cases. In the last example, the timing difference can be seen.

- (3) Consider $R = \mathbb{Q}[x, y, z, t, v, a_1, \dots, a_5]$,
 $I = (v^4 x^{12} z^{73}, v^{87} t^{21} y^{13}, x^{43} y^{18} z^{72} t^{28}, vxy, vyz, vzt, vtx, a_1^{7000}, a_2^{413}),$
 $J = (v^5 x^{13} z^{74}, v^{88} t^{22} y^{14}, x^{44} y^{19} z^{73} t^{29}, v^2 x^2 y^2, v^2 y^2 z^2, v^2 z^2 t^2, v^2 t^2 x^2).$
Timing for depth I/J computation: 16m 11s.
Timing for depth $\overline{I/J}$ computation: 11m.

2. APPENDIX

We sketch the simple idea of the algorithm which computes the canonical form of a monomial ideal I . This can easily be extended to compute the canonical form of I/J by simply applying it for $G(I) \cup G(J)$ and afterwards extracting the generators corresponding to I and J . This was used in Example 1.13.

The algorithm is based on Remark 1.4: for each variable x_i we build the list **gp** in which we save the pair (g, p) , where p is chosen such that x_i^p enters the g -generator of the monomial ideal I . This list will be sorted by the powers p as in the following example

Example 2.1. Consider the ideal $I := (x^{13}, x^4 y^7, y^7 z^{10}) \subset \mathbb{Q}[x, y, z]$. Then for each variable we will obtain a different **gp** as shown below:

- For the first variable x , **gp** is equal to $\boxed{2 \mid 4 \mid 1 \mid 13}$. Therefore I is of type $(4, 13)$ with respect to x . Hence, in order to obtain the canonical form with respect to x , one has to divide the second generator by $x^{4-1} = x^3$ and the first generator by $x^{13-2} = x^{11}$. After these computation we will get $I_1 = (x^2, xy^7, y^7 z^{10})$. Note that I_1 is in the canonical form w.r.t. x .
- For the second variable y , **gp** is equal to $\boxed{3 \mid 7 \mid 2 \mid 7}$. Similar as above, one has to divide the second and the third generator by y^6 , and hence it results $I_2 = (x^2, xy, yz^{10})$. Again, I_2 is in the canonical form w.r.t. y and x .
- For the last variable z , **gp** is equal to $\boxed{3 \mid 10}$. We divide the third generator of I_2 by z^9 and we get our final result $I' = (x^2, xy, yz)$, which is in the canonical form with respect to all variables.

Based on the above idea, we construct two procedures: **putIn** and **canonical** – the first one constructing the list **gp**, and the second one computing the canonical form of a monomial ideal. The proof of correctness and termination is trivial. The procedures were written in the SINGULAR language.

```
proc putIn(intvec v, int power, int nrgen)
{
  if(size(v) == 1)
  {
    v[1] = nrgen;
  }
}
```

```

        v[2] = power;
        return(v);
    }
    int i,j;
    if(power <= v[2])
    {
        for(j = size(v)+2; j >=3; j--)
        {
            v[j] = v[j-2];
        }
        v[1] = nrgen;
        v[2] = power;
        return(v);
    }
    if(power >= v[size(v)])
    {
        v[size(v)+1] = nrgen;
        v[size(v)+1] = power;
        return(v);
    }
    for(j = size(v) + 2; (j>=4) && (power < v[j-2]); j = j-2)
    {
        v[j] = v[j-2];
        v[j-1] = v[j-3];
    }
    v[j] = power;
    v[j-1] = nrgen;
    return(v);
}

```

```

proc canonical(ideal I)
{
    int i,j,k;
    intvec gp;
    ideal m;
    intvec v;
    v = 0:nvars(basering);
    for(i = 1; i<=nvars(basering); i++)
    {
        gp = 0;
        v[i] = 1;
        for(j = 1; j<=size(I); j++)
        {
            if(deg(I[j],v) >= 1)

```

```

        {
            gp = putIn(gp,deg(I[j],v),j);
        }
    }
    k = 0;
    if(size(gp) == 2)
    {
        I[gp[1]] = I[gp[1]]/(var(i)^(gp[2]-1));
    }
    else
    {
        for(j = 1; j<=size(gp)-2;)
        {
            k++;
            I[gp[j]] = I[gp[j]]/(var(i)^(gp[j+1]-k));
            j = j+2;
            while((j<=size(gp)-2) && (gp[j-1] == gp[j+1]) )
            {
                I[gp[j]] = I[gp[j]]/(var(i)^(gp[j+1]-k));
                j = j + 2;
            }
        }
        if(j == size(gp)-1)
        {
            if(gp[j-1] == gp[j+1])
            {
                I[gp[j]] = I[gp[j]]/(var(i)^(gp[j+1]-k));
            }
            else
            {
                k++;
                I[gp[j]] = I[gp[j]]/(var(i)^(gp[j+1]-k));
            }
        }
    }
    v[i] = 0;
}
return(I);
}

```

REFERENCES

- [1] J. Abbott, A. M. Bigatti: *CoCoALib: a C++ library for doing Computations in Commutative Algebra*, available at <http://cocoa.dima.unige.it/cocoalib>
- [2] M. Cimpoeas, *Stanley depth of complete intersection monomial ideals*, Bull. Math. Soc. Sc. Math. Roumanie **51**(99), 205-211, (2008).

- [3] W. Decker, G.-M. Greuel, G. Pfister, H. Schönemann: SINGULAR 3-1-6 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2012).
- [4] J. Herzog, M. Vladoiu, X. Zheng, *How to compute the Stanley depth of a monomial ideal*, J. Algebra, **322**, 3151-3169, (2009).
- [5] B. Ichim, L. Katthän, J. J. Moyano-Fernández, *The behaviour of Stanley depth under polarization*, arXiv:AC/1401.4309, (2014).
- [6] M. Ishaq, M. I. Qureshi, *Upper and lower bounds for the Stanley depth of certain classes of monomial ideals and their residue class rings*, Communications in Algebra, Volume 41, 1107-1116, (2013)
- [7] R. Okazaki, K. Yanagawa: *Alexander duality and Stanley depth of multigraded modules*, Journal of Algebra 340: 35-52, (2011).
- [8] A. Popescu: *An algorithm to compute the Hilbert depth*, to appear in Journal of Symbolic Computation, (2013), arXiv.org/AC/1307.6084v3
- [9] A. Popescu, D. Popescu, *Four generated, squarefree, monomial ideals*, to appear in Proceedings of the International Conference "Experimental and Theoretical Methods in Algebra, Geometry, and Topology, June 20-24, 2013", Editors Denis Ibadula, Willem Veys, Springer-Verlag, arxiv:AC/1309.4986v4, (2014)
- [10] D. Popescu, *An inequality between depth and Stanley depth*, Bull. Math. Soc. Sc. Math. Roumanie **52**(100), 377-382, (2009).
- [11] G. Rinaldo, *An algorithm to compute the Stanley depth of monomial ideals*, Le Matematiche, Vol. LXIII, 243-256, (2008).
- [12] R. P. Stanley, *Linear Diophantine equations and local cohomology*, Invent. Math. 68 175-193, (1982).
- [13] K. Yanagawa, *Sliding functor and polarization functor for multigraded modules*, Communications in Algebra, 40: 1151-1166, (2012).

ADRIAN POPESCU, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN, ERWIN-SCHRÖDINGER-STR., 67663 KAISERSLAUTERN, GERMANY
E-mail address: popescu@mathematik.uni-kl.de